

A BLOWUP CRITERIA ALONG MAXIMUM POINTS OF THE 3D-NAVIER-STOKES FLOW IN TERMS OF FUNCTION SPACES WITH VARIABLE GROWTH CONDITION

EIICHI NAKAI AND TSUYOSHI YONEDA

ABSTRACT. A blowup criteria along maximum point of the 3D-Navier-Stokes flow in terms of function spaces with variable growth condition is constructed. This criterion is different from the Beale-Kato-Majda type and Constantin-Fefferman type criterion. If geometric behavior of the velocity vector field near the maximum point has a kind of symmetry up to a possible blowup time, then the solution can be extended to be the strong solution beyond the possible blowup time.

Key words: blowup criterion, 3D Navier-Stokes equation, Campanato spaces with variable growth condition.

AMS Subject Classification (2010): 35Q30, 76D03, 76D05, 46E35

1. INTRODUCTION

In this paper we construct a blowup criteria along maximum points of the 3D-Navier-Stokes flow in terms of function spaces with variable growth condition. The Navier-Stokes equation is expressed as

$$(1.1) \quad \begin{cases} \partial_t v + (v \cdot \nabla)v - \Delta v + \nabla p = 0 & \text{in } \mathbb{R}^3 \times [0, T), \\ \nabla \cdot v = 0 & \text{in } \mathbb{R}^3 \times [0, T), \\ v_0 = v|_{t=0} & \text{in } \mathbb{R}^3, \end{cases}$$

where v is a vector field representing velocity of the fluid, and p is the pressure. The most significant blowup criterion must be the Beale-Kato-Majda criterion [1]. The Beale-Kato-Majda criterion is as follows:

Theorem 1.1. *Let $s > 1/2$, and let $v_0 \in H^s$ with $\operatorname{div} v_0 = 0$ in distribution sense. Suppose that v is a strong solution of (1.1). If*

$$(1.2) \quad \int_0^T \|\operatorname{curl} v(t)\|_\infty dt < \infty,$$

then v can be extended to the strong solution up to some T' with $T' > T$.

This blowup criterion was further improved by Giga [9], Kozono and Taniuchi [11], the authors [20], etc. On the other hand, Constantin and Fefferman [5] (see also [6]) took into account geometric structure of the vortex stretching term in the vorticity equations

to get another kind of blowup condition. They imposed vortex direction condition to the high vorticity part. This criterion was also further improved by, for example, Deng, Hou and Yu [7]. These two separate forms of criteria controlling the blow-up by magnitude and the direction of the vorticity respectively are interpolated by Chae [3]. For the detail of the blowup problem of the Navier-Stokes equation, see Fefferman [8] for example.

In this paper, we give a different type of blowup criterion from them. We focus on a geometric behavior of the velocity vector field near the each maximum points. In order to state our blowup criterion, we need to give several definitions.

Let us denote a maximum point of $|v|$ at a time t as $x_M = x_{M(t)} \in \mathbb{R}^3$ (if there are several maximum points at a time t , then we choose one maximum point. We sometimes abbreviate the time t). We use rotation and transformation and bring a maximum point to the origin and its direction parallel to x_3 -axis. Then we decompose v into two parts: symmetric flow part and its remainder. In this paper we prove that, if the remainder part is small, then the solution never blowup.

Let us explain precisely. We denote the unit tangent vector as

$$\tau(x_M) = \tau(x_{M(t)}) = (v/|v|)(x_{M(t)}, t),$$

and we choose unit normal vectors $n_1(x_M)$ and $n_2(x_M)$ as

$$\tau(x_M) \cdot n_1(x_M) = \tau(x_M) \cdot n_2(x_M) = n_1(x_M) \cdot n_2(x_M) = 0.$$

Note that n_1 and n_2 are not uniquely determined. We now construct a Cartesian coordinate system with a new y_1 -axis to be the straight line which passes through the maximum point and is parallel to n_1 , and a new y_2 -axis to be the straight line which passes through the maximum point and is parallel to n_2 . We set y_3 -axis by τ in the same process. Here we fix the maximum point $x_M = x_{M(t_*)}$ at $t = t_*$ for some time. Then v can be expressed as

$$(1.3) \quad v(x, t) = \tilde{u}_1(x, t)n_1(x_{M(t_*)}) + \tilde{u}_2(x, t)n_2(x_{M(t_*)}) + \tilde{u}_3(x, t)\tau(x_{M(t_*)}),$$

with $\tilde{u} = (\tilde{u}_1, \tilde{u}_2, \tilde{u}_3)$, where

$$\tilde{u}_1(x, t) = v(x, t) \cdot n_1(x_{M(t_*)}),$$

$$\tilde{u}_2(x, t) = v(x, t) \cdot n_2(x_{M(t_*)}),$$

$$\tilde{u}_3(x, t) = v(x, t) \cdot \tau(x_{M(t_*)}).$$

Let $y = (y_1, y_2, y_3)$ be the coordinate representation of the point x in the coordinate system based at the maximum point which is specified by the orthogonal frame $\{n_1, n_2, \tau\}$. That is, the point $x \in \mathbb{R}^3$ can be realized as $x = x_M + n_1(x_M)y_1 + n_2(x_M)y_2 + \tau(x_M)y_3$ with

$x_M = x_{M(t_*)}$. Then we can rewrite $\tilde{u}(x) = \tilde{u}(x, t)$ to $u(y) = u(y, t) = u_{M(t_*)}(y, t)$ as

$$\begin{aligned} u_1(y) &= u_1(y, t) = \tilde{u}_1(x_M + n_1(x_M)y_1 + n_2(x_M)y_2 + \tau(x_M)y_3, t), \\ u_2(y) &= u_2(y, t) = \tilde{u}_2(x_M + n_1(x_M)y_1 + n_2(x_M)y_2 + \tau(x_M)y_3, t), \\ u_3(y) &= u_3(y, t) = \tilde{u}_3(x_M + n_1(x_M)y_1 + n_2(x_M)y_2 + \tau(x_M)y_3, t). \end{aligned}$$

In this case $u_1(0, t_*) = u_2(0, t_*) = 0$ and $u_3(0, t_*) = |v(x_{M(t_*)}, t_*)|$.

Since the Navier-Stokes equation is rotation and translation invariant, u also satisfies the Navier-Stokes equation (1.1) in y -valuable. Then ∇p , in y -valuable, can be expressed as

$$\nabla p = \sum_{i,j=1}^3 R_i R_j \nabla (u_i u_j),$$

where R_j ($j = 1, 2, 3$) are the Riesz transforms. We decompose u into two parts; symmetric flow part U and its remainder part r :

$$u = U + r.$$

The symmetric flow part U can be defined as follows:

Definition 1.1. We say U is a symmetric flow if U satisfies

$$\begin{cases} U_1(y_1, y_2, y_3) = -U_1(y_1, y_2, -y_3), \\ U_2(y_1, y_2, y_3) = -U_2(y_1, y_2, -y_3), \\ U_3(y_1, y_2, y_3) = U_3(y_1, y_2, -y_3). \end{cases}$$

We see that the symmetric flow cannot create large gradient of the pressure. Actually, a basic calculation shows that

$$(1.4) \quad \sum_{i,j=1}^3 R_i R_j \partial_3 (U_i U_j)|_{y=0} = 0,$$

since, if f is even (odd) with respect to y_3 , then $R_1 f$ and $R_2 f$ are also even (odd) with respect to y_3 , but $R_3 f$ is odd (even) with respect to y_3 . Thus we need to see the remainder part r , namely, we have the following pressure formula:

$$(1.5) \quad \partial_3 p|_{y=0} = \sum_{i,j=1}^3 R_i R_j \partial_3 (r_i U_j + U_i r_j + r_i r_j)|_{y=0}.$$

In this paper, using the above formula, we construct a different type (from Beale-Kato-Majda type and Constantin-Fefferman type) of blowup criterion. We measure symmetry of the flow near each maximum points by controlling the remainder part r . In order to obtain a reasonable blowup condition from (1.5), we need two function spaces

$V = (V, \|\cdot\|_V)$ and $W = (W, \|\cdot\|_W)$ on $\mathbb{R}^3$ such that

$$(1.6) \quad |f(0)| \leq \|f\|_W,$$

$$(1.7) \quad \|R_i R_j f\|_W \leq C \|f\|_W,$$

$$(1.8) \quad \|fg\|_W \leq C \|f\|_V \|g\|_V.$$

That is, we need some smoothness condition at the origin for functions in W , the boundedness of Riesz transforms on W and the boundedness of pointwise multiplication operator as $V \times V \rightarrow W$. Moreover, it is known that there exist positive constants R and C such that

$$(1.9) \quad |v(x, t)| \leq C/|x| \quad \text{for } |x| > R,$$

where R and C are independent of $t \in [0, T]$. This is due to Corollary 1 in [2] (we use the partial regularity result to the decay). See also Section 1 in [4]. We need to take the decay condition (1.9) into account to construct V . In these points of view, we use Campanato spaces with variable growth condition. We discuss these function spaces in Sections 3–6. The following definition is the key in this paper.

Definition 1.2. We say “ v is no local collapsing (of its symmetricity near each maximum points)” with respect to the function space V , if there exist constants $C > 0$ and $\alpha < 2$ such that, for each fixed $x_{M(t_*)}$ at $t_* \in [0, T]$, $u = u_{M(t_*)}$ has the following property:

$$\inf_{u=U+r} \left\{ \sum_{i,j} (\|\partial_3 r_i\|_V \|U_j\|_V + \|r_i\|_V \|\partial_3 U_j\|_V + \|r_i\|_V \|\partial_3 r_j\|_V) \Big|_{t=t_*} \right\} \leq C \frac{(T - t_*)^{-\alpha}}{u_3(0, t_*)},$$

where the infimum is taken over all decomposition $u = U + r$ with symmetric flow U .

Roughly saying, if $\|\partial_3 r_j\|_V$ and $\|r_j\|_V$ are sufficiently small compare to $\|\partial_3 U_j\|_V$ and $\|U_j\|_V$ (which means symmetric part is dominant), then v is no local collapsing.

The following is the main theorem.

Theorem 1.2 (Blowup criteria along maximum points). *Let function spaces V and W satisfy (1.6), (1.7) and (1.8). Let v_0 be any non zero, smooth, divergence-free vector field in Schwartz class, that is,*

$$|\partial_x^\alpha v_0(x)| \leq C_{\alpha,K} (1 + |x|)^{-K} \quad \text{in } \mathbb{R}^3$$

for any $\alpha \in \mathbb{Z}_+^3$ and any $K > 0$. Suppose that $v \in C^\infty([0, T] \times \mathbb{R}^3)$ is a unique smooth solution of (1.1) up to T . If v is no local collapsing with respect to V , then v can be extended to the strong solution up to some T' with $T' > T$.

In the next section we prove Theorem 1.2 by using the regularity criterion by [9]. We also give an example of function with no local collapsing which doesn't satisfy the Beale-Kato-Majda criterion. In Section 3 we define Campanato spaces with variable growth

condition which give concrete function spaces V and W satisfying (1.6), (1.7) and (1.8). Campanato spaces with variable growth condition were introduced by [19] to characterize the pointwise multipliers on BMO, and then they were investigated by [14, 15, 17, 18], etc. Roughly saying, the function spaces V and W are required to express C^α ($0 < \alpha < 1$) continuity near the origin and the decay condition (1.9) far from the origin. For these requirement, we can use Campanato spaces with variable growth condition. We state the boundedness of the Riesz transforms and the pointwise multiplication operator on these function spaces in Section 4 and Section 5, respectively. Finally, we show that Campanato spaces satisfy the conditions (1.6), (1.7) and (1.8) for some variable growth condition in Section 6.

2. PROOF OF THE MAIN THEOREM

In this section we give a proof of the main theorem. First we show a lemma.

Lemma 2.1. *Under the assumption of Theorem 1.2, for each fixed $x_{M(t_*)}$, the following inequalities hold:*

$$(2.1) \quad -(v \cdot \nabla p)(x_{M(t_*)}, t_*) \leq C(T - t_*)^{-\alpha},$$

$$(2.2) \quad (v \cdot \Delta v)(x_{M(t_*)}, t_*) \leq 0.$$

Proof. Using the derivative ∂_3 along τ direction, we have

$$-(v \cdot \nabla p)(x_{M(t_*)}, t_*) = -(u_3 \partial_3 p)(0, t_*),$$

since $u_1(0, t_*) = u_2(0, t_*) = 0$. Then, by (1.5), (1.6) and the definition of no local collapsingness, we get (2.1).

Next we show (2.2). To do this we prove

$$(u_3 \Delta u_3)(0, t_*) \leq 0,$$

where Δ is the Laplacian with respect to $y = (y_1, y_2, y_3)$. Since $y = 0$ is a maximum point, we see

$$\partial_j |u(y)| \Big|_{y=0} = 0 \quad \text{for } j = 1, 2, 3,$$

and

$$\partial_j^2 |u(y)| \Big|_{y=0} \leq 0 \quad \text{for } j = 1, 2, 3.$$

There are smooth functions θ_1 , θ_2 and θ_3 such that

$$\begin{aligned} u_1(y) &= |u(y)| \sin \theta_1(y), \\ u_2(y) &= |u(y)| \sin \theta_2(y), \\ u_3(y) &= |u(y)| \cos \theta_3(y) \end{aligned}$$

with $\theta_1(0) = \theta_2(0) = \theta_3(0) = 0$. A direct calculation yields

$$\begin{aligned}\partial_1 u_3(y) &= \partial_1 |u(y)| \cos \theta_3(y) - |u(y)| \sin \theta_3(y) \partial_1 \theta_3(y) \\ \partial_1^2 u_3(y) &= \partial_1^2 |u(y)| \cos \theta_3(y) - 2 \partial_1 |u(y)| \sin \theta_3(y) \partial_1 \theta_3(y) \\ &\quad - |u(y)| \cos \theta_3(y) (\partial_1 \theta_3(y))^2 - |u(y)| \sin \theta_3(y) \partial_1^2 \theta_3(y).\end{aligned}$$

Thus we have

$$\left. \partial_1^2 u_3(y) \right|_{y=0} = \partial_{y_1}^2 |u(y)| - |u(y)| (\partial_{y_1} \theta_3(y))^2 \Big|_{y=0} \leq 0.$$

Similar calculations to y_2 and y_3 directions, we have $(u_3 \Delta u_3)(0, t_*) \leq 0$. $\square$

Next we define “trajectory” $\gamma : [\tilde{t}, T) \rightarrow \mathbb{R}^3$ starting at a point $\tilde{x}$:

$$\partial_t \gamma(\tilde{x}, \tilde{t}; t) = v(\gamma(\tilde{x}, \tilde{t}; t), t) \quad \text{with} \quad \gamma(\tilde{x}, \tilde{t}; \tilde{t}) = \tilde{x}.$$

Then γ provides a diffeomorphism and the equation (1.1) can be rewritten as follows:

$$\partial_t \left(v(\gamma(\tilde{x}, \tilde{t}; t), t) \right) = (\Delta v - \nabla p)(\gamma(\tilde{x}, \tilde{t}; t), t) \quad (\tilde{t} < t < T)$$

with $\gamma(\tilde{x}, \tilde{t}; \tilde{t}) = \tilde{x} \in \mathbb{R}^3$. Since v is bounded for fixed $t \in [0, T)$, we can define $X(t) \subset \mathbb{R}^3$ as the set of all maximum points of $|v(\cdot, t)|$ at a time $t \in [0, T)$, namely,

$$|v(x, t)| = \sup_{\xi \in \mathbb{R}^3} |v(\xi, t)| \text{ for } x \in X(t) \quad \text{and} \quad |v(x, t)| < \sup_{\xi \in \mathbb{R}^3} |v(\xi, t)| \text{ for } x \notin X(t).$$

By (1.9), $X(t)$ is a bounded set uniformly in t in a possible blowup scenario. Let $B(x, r)$ is a ball with radius r and centered at x . For any $r > 0$, we see that there is a barrier function $\beta(t) > 0$ such that

$$|v(x, t)| + \beta(t) < \sup_{\xi \in \mathbb{R}^3} |v(\xi, t)| \quad \text{for} \quad x \notin \cup_{\xi \in X(t)} B(\xi, r).$$

Then, using Lemma 2.1 and the smoothness of the solution, we get the following:

Proposition 2.2. *Under the assumption of Theorem 1.2, for any $\delta > 0$ and $t_* \in [0, T)$, there exists a time interval $[t_*, t'_*) \subset [0, T)$ and a radius r_* such that the following two properties hold for all $t' \in [t_*, t'_*)$:*

- $\cup_{\xi \in X(t_*)} B(\xi, r_*) \Subset \Omega(t')$, where

$$(2.3) \quad \Omega(t') := \left\{ x \in \mathbb{R}^3 : (\Delta v \cdot v)(\gamma(x, t_*; t'), t') \leq \delta, \right. \\ \left. (-\nabla p \cdot v)(\gamma(x, t_*; t'), t') \leq \delta + C(T - t')^{-\alpha} \right\},$$

- $|v(\gamma(x, t_*; t'), t')|^2 < \sup_{\xi \in \mathbb{R}^3} |v(\xi, t_*)|^2$ for $x \in (\cup_{\xi \in X(t_*)} B(\xi, r_*))^c$.

Proof of Theorem 1.2. Note that the open interval $(0, T)$ is covered by the collection $\{(t_*, t'_*)\}_{t_* \in [0, T]}$ of the open intervals such that the interval $[t_*, t'_*)$ is as in Proposition 2.2 for $t_* \in [0, T)$. Since $(0, T)$ is a Lindelöf space, we can choose a sequence of the time intervals $[t_j, t'_j)$, $j = 0, 1, 2, \dots$ (finite or infinite), such that $(0, T) = \cup_j (t_j, t'_j)$, and that $[t_j, t'_j)$ and r_j satisfy the properties of Proposition 2.2 for $t_j \in [0, T)$. We may assume that

$$0 = t_0 < t_1 < t_2 < \dots, \quad t_{j+1} < t'_j, \quad j = 0, 1, \dots$$

For $t \in [t_0, t'_0)$ and $x \in \cup_{\xi \in X(t_0)} B(\xi, r_0)$, from the first property in Proposition 2.2 it follows that

$$\begin{aligned} & |v(\gamma(x, t_0; t), t)|^2 \\ &= \int_{t_0}^t \partial_{t'} |v(\gamma(x, t_0; t'), t')|^2 dt' + |v(x, t_0)|^2 \\ &= 2 \int_{t_0}^t \partial_{t'} v \cdot v dt' + |v(x, 0)|^2 \\ &= 2 \int_{t_0}^t (\Delta v \cdot v - \nabla p \cdot v) dt' + |v(x, t_0)|^2 \\ &\leq 2 \left(2\delta(t - t_0) + C \int_{t_0}^t (T - t')^{-\alpha} dt' \right) + \sup_{\xi \in \mathbb{R}^3} |v(\xi, t_0)|^2. \end{aligned}$$

The case $x \in (\cup_{\xi \in X(t_0)} B(\xi, r_0))^c$ is straightforward by the second property in Proposition 2.2. Then we have

$$|v(z, t)|^2 \leq 2 \left(2\delta(t - t_0) + C \int_{t_0}^t (T - t')^{-\alpha} dt' \right) + \sup_{\xi \in \mathbb{R}^3} |v(\xi, t_0)|^2.$$

for all $t \in [t_0, t'_0)$ and all $z \in \mathbb{R}^3$ with $z = \gamma(x, t_0; t)$, since γ gives a diffeomorphism. Repeating the above argument infinite times, and we finally have

$$|v(x, t)|^2 \leq 2 \left(2\delta t + C \int_0^t (T - t')^{-\alpha} dt' \right) + \sup_{\xi \in \mathbb{R}^3} |v(\xi, 0)|^2$$

for all $t \in [0, T)$ and all $x \in \mathbb{R}^3$. This implies

$$\|v\|_{L^2(0, T; L^\infty(\mathbb{R}^3))} < \infty.$$

Due to the classical regularity criterion (see [9] for example), we see that the solution never blowup. $\square$

Remark 2.1. We can construct a function u which satisfy both Definition 1.2 and

$$\int_0^T \|\operatorname{curl} u(t)\|_\infty = \infty \quad (\text{the Beale-Kato-Majda criterion})$$

(in this remark, u is nothing to do with the Navier-Stokes solution, we just regard u as a time dependent vector field). If $\theta_j(y) = \theta_j(-y)$ ($j = 1, 2, 3$, even angular), we see that

$\partial_3 u_1(y) - \partial_1 u_3(y)|_{y=0}$ is allowed to be arbitrary large. In fact,

$$\begin{aligned}\partial_1 u_3(y) &= (\partial_1 |u(y)|) \cos \theta_3(y) - |u(y)| \sin \theta_3(y) \partial_1 \theta_3(y), \\ \partial_3 u_1(y) &= (\partial_3 |u(y)|) \sin \theta_1(y) + |u(y)| \cos \theta_1(y) \partial_3 \theta_1(y)\end{aligned}$$

and then

$$\left. \partial_3 u_1(y) - \partial_1 u_3(y) \right|_{y=0} = |u(y)| \partial_3 \theta_1(y) \Big|_{y=0}.$$

Since $\partial_3 \theta_1(0)$ can be taken arbitrary large for each $t > 0$, we can construct the desired function u . Note that since $\theta_j(y)$ ($j = 1, 2, 3$) are even angular, u is symmetric flow (see Definition 1.1).

3. CAMPANATO SPACES WITH VARIABLE GROWTH CONDITION

In this section we define Campanato spaces $\mathcal{L}_{p,\phi}^\natural$ with variable growth condition. We state basic properties of the function spaces $\mathcal{L}_{p,\phi}^\natural$. To do this we also define Morrey spaces and Hölder spaces with variable growth condition.

Let $\mathbb{R}^n$ be the n -dimensional Euclidean space. We denote by $B(x, r)$ the open ball centered at $x \in \mathbb{R}^n$ and of radius r , that is,

$$B(x, r) = \{y \in \mathbb{R}^n : |y - x| < r\}.$$

For a measurable set $G \subset \mathbb{R}^n$, we denote by $|G|$ and χ_G the Lebesgue measure of G and the characteristic function of G , respectively.

We consider variable growth functions $\phi : \mathbb{R}^n \times (0, \infty) \rightarrow (0, \infty)$. For a ball $B = B(x, r)$, write $\phi(B)$ in place of $\phi(x, r)$. For a function $f \in L_{\text{loc}}^1(\mathbb{R}^n)$ and for a ball B , let

$$f_B = |B|^{-1} \int_B f(x) dx.$$

Then we define Campanato spaces $\mathcal{L}_{p,\phi}(\mathbb{R}^n)$ and $\mathcal{L}_{p,\phi}^\natural(\mathbb{R}^n)$, Morrey spaces $L_{p,\phi}(\mathbb{R}^n)$, and Hölder spaces $\Lambda_\phi(\mathbb{R}^n)$ and $\Lambda_\phi^\natural(\mathbb{R}^n)$ with variable growth functions ϕ as the following:

Definition 3.1. For $1 \leq p < \infty$ and $\phi : \mathbb{R}^n \times (0, \infty) \rightarrow (0, \infty)$, function spaces $\mathcal{L}_{p,\phi}(\mathbb{R}^n)$, $\mathcal{L}_{p,\phi}^\natural(\mathbb{R}^n)$, $L_{p,\phi}(\mathbb{R}^n)$, $\Lambda_\phi(\mathbb{R}^n)$, $\Lambda_\phi^\natural(\mathbb{R}^n)$ are the set of all functions f such that

$$\begin{aligned}\|f\|_{\mathcal{L}_{p,\phi}} &= \sup_B \frac{1}{\phi(B)} \left(\frac{1}{|B|} \int_B |f(x) - f_B|^p dx \right)^{1/p} < \infty, \\ \|f\|_{\mathcal{L}_{p,\phi}^\natural} &= \|f\|_{\mathcal{L}_{p,\phi}} + |f_{B(0,1)}| < \infty, \\ \|f\|_{L_{p,\phi}} &= \sup_B \frac{1}{\phi(B)} \left(\frac{1}{|B|} \int_B |f(x)|^p dx \right)^{1/p} < \infty, \\ \|f\|_{\Lambda_\phi} &= \sup_{x,y \in \mathbb{R}^n, x \neq y} \frac{2|f(x) - f(y)|}{\phi(x, |x - y|) + \phi(y, |y - x|)} < \infty, \\ \|f\|_{\Lambda_\phi^\natural} &= \|f\|_{\Lambda_\phi} + |f(0)| < \infty,\end{aligned}$$

respectively.

We regard $\mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$ and $L_{p,\phi}(\mathbb{R}^n)$ as spaces of functions modulo null-functions, $\mathcal{L}_{p,\phi}(\mathbb{R}^n)$ as spaces of functions modulo null-functions and constant functions, $\Lambda_{\phi}^{\natural}(\mathbb{R}^n)$ as a space of functions defined at all $x \in \mathbb{R}^n$, and $\Lambda_{\phi}(\mathbb{R}^n)$ as a space of functions defined at all $x \in \mathbb{R}^n$ modulo constant functions. Then these five functionals are norms and thereby these spaces are all Banach spaces.

In order to apply $\mathcal{L}_{p,\phi}^{\natural}$ to the blowup criterion (more precisely, in order to find specific function spaces V and W satisfying (1.6), (1.7) and (1.8)), we state several properties of these function spaces and relation between ϕ and the function spaces. For two variable growth functions ϕ_1 and ϕ_2 , we write $\phi_1 \sim \phi_2$ if there exists a positive constant C such that

$$C^{-1}\phi_1(B) \leq \phi_2(B) \leq C\phi_1(B) \quad \text{for all balls } B.$$

In this case, two spaces defined by ϕ_1 and by ϕ_2 coincide with equivalent norms. If $p = 1$ and $\phi \equiv 1$, then $\mathcal{L}_{p,\phi}(\mathbb{R}^n)$ is the usual $\text{BMO}(\mathbb{R}^n)$. For $\phi(x, r) = r^{\alpha}$, $0 < \alpha \leq 1$, we denote $\Lambda_{r^{\alpha}}(\mathbb{R}^n)$ and $\Lambda_{r^{\alpha}}^{\natural}(\mathbb{R}^n)$ by $\text{Lip}_{\alpha}(\mathbb{R}^n)$ and $\text{Lip}_{\alpha}^{\natural}(\mathbb{R}^n)$, respectively. In this case,

$$\|f\|_{\text{Lip}_{\alpha}} = \sup_{x,y \in \mathbb{R}^n, x \neq y} \frac{|f(x) - f(y)|}{|x - y|^{\alpha}} \quad \text{and} \quad \|f\|_{\text{Lip}_{\alpha}^{\natural}} = \|f\|_{\text{Lip}_{\alpha}} + |f(0)|.$$

If $\phi(x, r) = \min(r^{\alpha}, 1)$, $0 < \alpha \leq 1$, then

$$\|f\|_{\Lambda_{\phi}^{\natural}} \sim \|f\|_{\text{Lip}_{\alpha}} + \|f\|_{L^{\infty}}.$$

From the definition it follows that

$$\|f\|_{\mathcal{L}_{p,\phi}} \leq 2\|f\|_{L_{p,\phi}}, \quad \|f\|_{\mathcal{L}_{p,\phi}^{\natural}} \leq (2 + \phi(0, 1))\|f\|_{L_{p,\phi}}.$$

If $\phi(B) = |B|^{-1/p}$ for all balls B , then

$$\|f\|_{L_{p,\phi}} = \|f\|_{L^p}.$$

We consider the following conditions on variable growth function ϕ :

$$(3.1) \quad \frac{1}{A_1} \leq \frac{\phi(x, s)}{\phi(x, r)} \leq A_1, \quad \frac{1}{2} \leq \frac{s}{r} \leq 2,$$

$$(3.2) \quad \frac{1}{A_2} \leq \frac{\phi(x, r)}{\phi(y, r)} \leq A_2, \quad d(x, y) \leq r,$$

$$(3.3) \quad \phi(x, r) \leq A_3\phi(x, s), \quad 0 < r < s < \infty,$$

where A_i , $i = 1, 2, 3$, are positive constants independent of $x, y \in \mathbb{R}^n$, $r, s > 0$. Note that (3.2) and (3.3) imply that there exists a positive constant C such that

$$\phi(x, r) \leq C\phi(y, s) \quad \text{for } B(x, r) \subset B(y, s),$$

where the constant C is independent of balls $B(x, r)$ and $B(y, s)$.

The following three theorems are known:

Theorem 3.1 ([16]). *If ϕ satisfies (3.1), (3.2) and (3.3), then, for every $1 \leq p < \infty$, $\mathcal{L}_{p,\phi}(\mathbb{R}^n) = \mathcal{L}_{1,\phi}(\mathbb{R}^n)$ and $\mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n) = \mathcal{L}_{1,\phi}^{\natural}(\mathbb{R}^n)$ with equivalent norms, respectively.*

Theorem 3.2 ([15]). *If ϕ satisfies (3.1), (3.2), (3.3), and there exists a positive constant C such that*

$$(3.4) \quad \int_0^r \frac{\phi(x,t)}{t} dt \leq C\phi(x,r), \quad x \in \mathbb{R}^n, \quad r > 0,$$

then, for every $1 \leq p < \infty$, each element in $\mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$ can be regarded as a continuous function, (that is, each element is equivalent to a continuous function modulo null-functions) and $\mathcal{L}_{p,\phi}(\mathbb{R}^n) = \Lambda_{\phi}(\mathbb{R}^n)$ and $\mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n) = \Lambda_{\phi}^{\natural}(\mathbb{R}^n)$ with equivalent norms, respectively. In particular, if $\phi(x,r) = r^{\alpha}$, $0 < \alpha \leq 1$, then, for every $1 \leq p < \infty$, $\mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n) = \text{Lip}_{\alpha}^{\natural}(\mathbb{R}^n)$ and $\mathcal{L}_{p,\phi}(\mathbb{R}^n) = \text{Lip}_{\alpha}(\mathbb{R}^n)$ with equivalent norms, respectively.

Theorem 3.3 ([15]). *Let $1 \leq p < \infty$. If ϕ satisfies (3.1), (3.2), and there exists a positive constant C such that*

$$(3.5) \quad \int_r^{\infty} \frac{\phi(x,t)}{t} dt \leq C\phi(x,r), \quad x \in \mathbb{R}^n, \quad r > 0,$$

then, for $f \in \mathcal{L}_{p,\phi}(\mathbb{R}^n)$, the limit $\sigma(f) = \lim_{r \rightarrow \infty} f_{B(0,r)}$ exists and

$$\|f\|_{\mathcal{L}_{p,\phi}} \sim \|f - \sigma(f)\|_{L_{p,\phi}}.$$

That is, the mapping $f \mapsto f - \sigma(f)$ is bijective and bicontinuous from $\mathcal{L}_{p,\phi}(\mathbb{R}^n)$ (modulo constants) to $L_{p,\phi}(\mathbb{R}^n)$.

Remark 3.1. If $\int_1^{\infty} \phi(0,t)/t dt < \infty$, then $\phi(0,r) \rightarrow 0$ as $r \rightarrow \infty$. Then, for $f \in L_{p,\phi}(\mathbb{R}^n)$, we have

$$|\sigma(f)| = \lim_{r \rightarrow \infty} |f_{B(0,r)}| \leq \lim_{r \rightarrow \infty} \phi(0,r) \|f\|_{L_{p,\phi}} \rightarrow 0 \quad \text{as } r \rightarrow \infty.$$

That is, $\sigma(f) = 0$.

For a ball $B_* \subset \mathbb{R}^n$ and $0 < \alpha \leq 1$, let

$$\|f\|_{\text{Lip}_{\alpha}(B_*)} = \sup_{x,y \in B_*, x \neq y} \frac{|f(x) - f(y)|}{|x - y|^{\alpha}}.$$

We also conclude the following:

Proposition 3.4. *Let $1 \leq p < \infty$ and $0 < \alpha \leq 1$. Assume that, for a ball B_* ,*

$$(3.6) \quad \phi(x,r) = r^{\alpha} \quad \text{for all balls } B(x,r) \subset B_*.$$

Then each element f in $\mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$ can be regarded as a continuous function on the ball B_* , and, there exists a positive constant C such that

$$\|f\|_{\text{Lip}_{\alpha}(B_*)} \leq C \|f\|_{\mathcal{L}_{p,\phi}},$$

where C is dependent only on n and α . In particular, if (3.6) holds for $B_* = B(0, 1)$, then each $f \in \mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$ is α -Lipschitz continuous near the origin and

$$\|f\|_{\mathcal{L}_{p,\phi}^{\natural}} \sim \|f\|_{\mathcal{L}_{p,\phi}} + |f(0)|.$$

Proof. It is known that, if ϕ satisfies (3.1), then

$$(3.7) \quad |f_{B(x,r_1)} - f_{B(x,r_2)}| \leq C \int_{r_1}^{2r_2} \frac{\phi(x,t)}{t} dt \|f\|_{\mathcal{L}_{p,\phi}} \quad \text{for } x \in \mathbb{R}^n, r_1 < r_2,$$

where C is dependent only on n , see [12, Lemma 2.4]. Hence we have that, if $B(x, r)$, $B(y, r) \subset B_*$, then

$$|f_{B(x,r)} - f_{B(y,r)}| \leq C \int_r^{2r+|x-y|} \frac{t^\alpha}{t} dt \|f\|_{\mathcal{L}_{p,\phi}} \leq C_*(2r + |x - y|)^\alpha \|f\|_{\mathcal{L}_{p,\phi}},$$

since $B(x, r)$, $B(y, r) \subset B((x+y)/2, r + |x-y|/2)$, where C_* is dependent only on n and α . Letting $r \rightarrow 0$, we have

$$|f(x) - f(y)| \leq C_* |x - y|^\alpha \|f\|_{\mathcal{L}_{p,\phi}},$$

for almost every $x, y \in B_*$. In this case we can regard that f is a continuous function modulo null-functions and we have

$$\|f\|_{\text{Lip}_\alpha(B_*)} \leq C_* \|f\|_{\mathcal{L}_{p,\phi}}.$$

If $B_* = B(0, 1)$, then

$$|f_{B(0,r)} - f_{B(0,1)}| \leq C \int_r^2 \frac{t^\alpha}{t} dt \|f\|_{\mathcal{L}_{p,\phi}} \leq C \|f\|_{\mathcal{L}_{p,\phi}}.$$

Letting $r \rightarrow 0$, we have

$$|f(0) - f_{B(0,1)}| \leq C \|f\|_{\mathcal{L}_{p,\phi}}.$$

This shows that $\|f\|_{\mathcal{L}_{p,\phi}} + |f_{B(0,1)}| \sim \|f\|_{\mathcal{L}_{p,\phi}} + |f(0)|$. $\square$

Proposition 3.5. *Let $1 \leq p < \infty$ and B_* be a ball such that $B(0, 1) \subset B_*$. Assume that there exists a positive constant A such that*

$$\phi(B) \leq A|B|^{-1/p} \quad \text{for all balls } B \subset B_*.$$

Then there exists a positive constant C such that

$$\left(\int_{B_*} |f(x)|^p dx \right)^{1/p} \leq C \|f\|_{\mathcal{L}_{p,\phi}^{\natural}}.$$

for all $f \in \mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$, where C is dependent only on A , n and p .

Proof. Let $B_* = B(x_*, r_*)$. Using (3.7), we have

$$|f_{B(0,1)} - f_{B(x_*, r_*)}| \leq C \int_1^{2r_*} \frac{At^{-n/p}}{t} dt \|f\|_{\mathcal{L}_{p,\phi}} \leq C_* \|f\|_{\mathcal{L}_{p,\phi}},$$

where C_* is dependent only on A , n and p . Then

$$\begin{aligned} \left(\int_{B_*} |f(x)|^p dx \right)^{1/p} &\leq \left(\int_{B_*} |f(x) - f_{B_*}|^p dx \right)^{1/p} + |f_{B(0,1)} - f_{B(x_*, r_*)}| + |f_{B(0,1)}| \\ &\leq (A + C_*) \|f\|_{\mathcal{L}_{p,\phi}} + |f_{B(0,1)}| \\ &\leq (A + C_* + 1) \|f\|_{\mathcal{L}_{p,\phi}^{\natural}}. \end{aligned}$$

This shows the conclusion. $\square$

4. SINGULAR INTEGRAL OPERATORS

In this section we consider the singular integral theory to show the boundedness of Riesz transforms in Campanato spaces with variable growth condition. We denote by $L_c^p(\mathbb{R}^n)$ the set of all $f \in L^p(\mathbb{R}^n)$ with compact support. Let $0 < \kappa \leq 1$. We shall consider a singular integral operator T with measurable kernel K on $\mathbb{R}^n \times \mathbb{R}^n$ satisfying the following properties:

$$(4.1) \quad |K(x, y)| \leq \frac{C}{|x - y|^n} \quad \text{for } x \neq y,$$

$$(4.2) \quad |K(x, y) - K(z, y)| + |K(y, x) - K(y, z)| \leq \frac{C}{|x - y|^n} \left(\frac{|x - z|}{|x - y|} \right)^\kappa$$

for $|x - y| \geq 2|x - z|$,

$$(4.3) \quad \int_{r \leq |x-y| < R} K(x, y) dy = \int_{r \leq |x-y| < R} K(y, x) dy = 0$$

for $0 < r < R < \infty$ and $x \in \mathbb{R}^n$,

where C is a positive constant independent of $x, y, z \in \mathbb{R}^n$. For $\eta > 0$, let

$$T_\eta f(x) = \int_{|x-y| \geq \eta} K(x, y) f(y) dy.$$

Then $T_\eta f(x)$ is well defined for $f \in L_c^p(\mathbb{R}^n)$, $1 < p < \infty$. We assume that, for all $1 < p < \infty$, there exists positive constant C_p independently $\eta > 0$ such that,

$$\|T_\eta f\|_{L^p} \leq C_p \|f\|_{L^p} \quad \text{for } f \in L_c^p(\mathbb{R}^n),$$

and $T_\eta f$ converges to Tf in $L^p(\mathbb{R}^n)$ as $\eta \rightarrow 0$. By this assumption, the operator T can be extended as a continuous linear operator on $L^p(\mathbb{R}^n)$. We shall say the operator T satisfying the above conditions is a singular integral operator of type κ . For example, Riesz transforms are singular integral operators of type 1.

Now, to define T for functions $f \in \mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$, we first define the modified version of T_η by

$$(4.4) \quad \tilde{T}_\eta f(x) = \int_{|x-y| \geq \eta} f(y) [K(x, y) - K(0, y)(1 - \chi_{B(0,1)}(y))] dy.$$

Then we can show that the integral in the definition above converges absolutely for each x and that $\tilde{T}_\eta f$ converges in $L^p(B)$ as $\eta \rightarrow 0$ for each ball B . We denote the limit by $\tilde{T}f$. If both $\tilde{T}f$ and Tf are well defined, then the difference is a constant.

We can show the following results. Theorem 4.1 is an extension of [17, Theorem 4.1] and Theorem 4.3 is an extension of [13, Theorem 2]. The proofs are almost the same.

Theorem 4.1. *Let $0 < \kappa \leq 1$ and $1 < p < \infty$. Assume that ϕ and ψ satisfy (3.1) and that there exists a positive constant A such that, for all $x \in \mathbb{R}^n$ and $r > 0$,*

$$(4.5) \quad r^\kappa \int_r^\infty \frac{\phi(x, t)}{t^{1+\kappa}} dt \leq A\psi(x, r).$$

If T is a singular integral operator of type κ , then $\tilde{T}$ is bounded from $\mathcal{L}_{p,\phi}(\mathbb{R}^n)$ to $\mathcal{L}_{p,\psi}(\mathbb{R}^n)$ and from $\mathcal{L}_{p,\phi}^\sharp(\mathbb{R}^n)$ to $\mathcal{L}_{p,\psi}^\sharp(\mathbb{R}^n)$, that is, there exists a positive constants C such that

$$\|\tilde{T}f\|_{\mathcal{L}_{p,\psi}} \leq C\|f\|_{\mathcal{L}_{p,\phi}}, \quad \|\tilde{T}f\|_{\mathcal{L}_{p,\psi}^\sharp} \leq C\|f\|_{\mathcal{L}_{p,\phi}^\sharp}.$$

Moreover, if ϕ and ψ satisfy (3.2) and (3.3) also, then $\tilde{T}$ is bounded from $\mathcal{L}_{1,\phi}^\sharp(\mathbb{R}^n)$ to $\mathcal{L}_{1,\psi}^\sharp(\mathbb{R}^n)$.

Corollary 4.2. *Under the assumption in Theorem 4.1, if ϕ and ψ satisfies (3.2), (3.3) and (3.4), then $\tilde{T}$ is bounded from $\Lambda_\phi(\mathbb{R}^n)$ to $\Lambda_\psi(\mathbb{R}^n)$ and from $\Lambda_\phi^\sharp(\mathbb{R}^n)$ to $\Lambda_\psi^\sharp(\mathbb{R}^n)$.*

For Morrey spaces $L_{p,\phi}(\mathbb{R}^n)$, we have the following.

Theorem 4.3. *Let $0 < \kappa \leq 1$ and $1 < p < \infty$. Assume that ϕ and ψ satisfy (3.1) and that there exists a positive constant A such that, for all $x \in \mathbb{R}^n$ and $r > 0$,*

$$\int_r^\infty \frac{\phi(x, t)}{t} dt \leq A\psi(x, r).$$

If T is a singular integral operator of type κ , then T is bounded from $L_{p,\phi}(\mathbb{R}^n)$ to $L_{p,\psi}(\mathbb{R}^n)$.

Now we state the boundedness of Riesz transforms. For f in Schwartz class, the Riesz transforms of f are defined by

$$R_j f(x) = c_n \lim_{\varepsilon \rightarrow 0} R_{j,\varepsilon} f(x), \quad j = 1, \dots, n,$$

where

$$R_{j,\varepsilon} f(x) = \int_{\mathbb{R}^n \setminus B(x,\varepsilon)} \frac{x_j - y_j}{|x - y|^{n+1}} f(y) dy, \quad c_n = \Gamma\left(\frac{n+1}{2}\right) \pi^{-\frac{n+1}{2}}.$$

Then it is known that there exists a positive constant C_p independently $\varepsilon > 0$ such that,

$$\|R_{j,\varepsilon} f\|_{L^p} \leq C_p \|f\|_{L^p} \quad \text{for } f \in L_c^p(\mathbb{R}^n),$$

and $R_{j,\varepsilon} f$ converges to $R_j f$ in $L^p(\mathbb{R}^n)$ as $\varepsilon \rightarrow 0$. That is, the operator R_j can be extended as a continuous linear operator on $L^p(\mathbb{R}^n)$. Hence, we can define a modified Riesz

transforms of f as

$$\tilde{R}_j f(x) = c_n \lim_{\varepsilon \rightarrow 0} \tilde{R}_{j,\varepsilon} f(x), \quad j = 1, \dots, n,$$

and

$$\tilde{R}_{j,\varepsilon} f(x) = \int_{\mathbb{R}^n \setminus B(x,\varepsilon)} \left(\frac{x_j - y_j}{|x - y|^{n+1}} - \frac{(-y_j)(1 - \chi_{B(0,1)}(y))}{|y|^{n+1}} \right) f(y) dy.$$

We note that, if both $R_j f$ and $\tilde{R}_j f$ are well defined on $\mathbb{R}^n$, then $R_j f - \tilde{R}_j f$ is a constant function. More precisely,

$$R_j f(x) - \tilde{R}_j f(x) = c_n \int_{\mathbb{R}^n} \frac{(-y_j)(1 - \chi_{B(0,1)}(y))}{|y|^{n+1}} f(y) dy.$$

Remark 4.1. If f is a constant function, then $\tilde{R}_j f = 0$. Actually, for $f \equiv 1$,

$$\begin{aligned} \tilde{R}_{j,\varepsilon} 1(x) &= \int_{\mathbb{R}^n \setminus B(x,\varepsilon)} \frac{(x_j - y_j) \chi_{B(x,1)}}{|x - y|^{n+1}} dy \\ &\quad + \int_{\mathbb{R}^n \setminus B(x,\varepsilon)} \left(\frac{(x_j - y_j)(1 - \chi_{B(x,1)})}{|x - y|^{n+1}} - \frac{(-y_j)(1 - \chi_{B(0,1)}(y))}{|y|^{n+1}} \right) dy \\ &= \int_{B(0,1) \setminus B(0,\varepsilon)} \frac{y_j}{|y|^{n+1}} dy + \int_{B(x,\varepsilon)} \frac{(-y_j)(1 - \chi_{B(0,1)}(y))}{|y|^{n+1}} dy \\ &= \int_{B(x,\varepsilon)} \frac{(-y_j)(1 - \chi_{B(0,1)}(y))}{|y|^{n+1}} dy \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

since

$$\int_{B(0,1) \setminus B(0,\varepsilon)} \frac{y_j}{|y|^{n+1}} dy = 0$$

and

$$\int_{\mathbb{R}^n} \left(\frac{(x_j - y_j)(1 - \chi_{B(x,1)})}{|x - y|^{n+1}} - \frac{(-y_j)(1 - \chi_{B(0,1)}(y))}{|y|^{n+1}} \right) dy = 0.$$

Hence $\tilde{R}_j 1(x) = 0$ for all $x \in \mathbb{R}^3$.

Theorem 4.4. Let $1 \leq p < \infty$ and ϕ satisfy (3.1) and

$$r \int_r^\infty \frac{\phi(x, t)}{t^2} dt \leq A\phi(x, r),$$

for all $x \in \mathbb{R}^n$ and $r > 0$. Assume that there exists a growth function $\tilde{\phi}$ such that $\phi \leq \tilde{\phi}$ and that $\tilde{\phi}$ satisfies (3.1), (3.2) and (3.5). If $f \in \mathcal{L}_{p,\phi}^\natural(\mathbb{R}^n)$ and $\sigma(f) = \lim_{r \rightarrow \infty} f_{B(0,r)} = 0$, then $R_j f$, $j = 1, 2, \dots, n$, are well defined, $\sigma(R_j f) = \lim_{r \rightarrow \infty} (R_j f)_{B(0,r)} = 0$, and

$$\|R_j f\|_{\mathcal{L}_{p,\phi}^\natural} \leq C \|f\|_{\mathcal{L}_{p,\phi}^\natural}, \quad j = 1, 2, \dots, n,$$

where C is a positive constant independent of f .

Proof. Let $f \in \mathcal{L}_{p,\phi}^\natural(\mathbb{R}^n)$ and $\sigma(f) = 0$. Then, by Theorem 3.3,

$$\|f\|_{L_{p,\tilde{\phi}}} = \|f - \sigma(f)\|_{L_{p,\tilde{\phi}}} \sim \|f\|_{\mathcal{L}_{p,\tilde{\phi}}} \leq \|f\|_{\mathcal{L}_{p,\phi}} \leq \|f\|_{\mathcal{L}_{p,\phi}^\natural}.$$

By Theorems 4.3 $R_j f$ is well defined and

$$\|R_j f\|_{L_{p,\tilde{\phi}}} \leq C \|f\|_{L_{p,\tilde{\phi}}} \leq C \|f\|_{\mathcal{L}_{p,\phi}^{\natural}}.$$

This shows that $\sigma(R_j f) = 0$ by Remark 3.1 and

$$|(R_j f)_{B(0,1)}| \leq \left(\frac{1}{|B(0,1)|} \int_{B(0,1)} |R_j f(x)|^p dx \right)^{1/p} \leq \tilde{\phi}(0,1) \|R_j f\|_{L_{p,\tilde{\phi}}} \leq C \|f\|_{\mathcal{L}_{p,\phi}^{\natural}}.$$

Since $R_j f - \tilde{R}_j f$ is a constant, by Theorem 4.1, we have

$$\|R_j f\|_{\mathcal{L}_{p,\phi}} = \|\tilde{R}_j f\|_{\mathcal{L}_{p,\phi}} \leq C \|f\|_{\mathcal{L}_{p,\phi}} \leq C \|f\|_{\mathcal{L}_{p,\phi}^{\natural}}.$$

Therefore, we have $\|R_j f\|_{\mathcal{L}_{p,\phi}^{\natural}} \leq C \|f\|_{\mathcal{L}_{p,\phi}^{\natural}}$. $\square$

5. POINTWISE MULTIPLICATION

Let $L^0(\mathbb{R}^n)$ be the set of all measurable functions on $\mathbb{R}^n$. Let X_1 and X_2 be subspaces of $L^0(\mathbb{R}^n)$ and $g \in L^0(\mathbb{R}^n)$. We say that g is a pointwise multiplier from X_1 to X_2 if $fg \in X_2$ for all $f \in X_1$. We denote by $\text{PWM}(X_1, X_2)$ the set of all pointwise multipliers from X_1 to X_2 .

For $\phi : \mathbb{R}^n \times (0, \infty) \rightarrow (0, \infty)$, we define

$$(5.1) \quad \Phi^*(x, r) = \int_1^{\max(2, |x|, r)} \frac{\phi(0, t)}{t} dt,$$

$$(5.2) \quad \Phi^{**}(x, r) = \int_r^{\max(2, |x|, r)} \frac{\phi(x, t)}{t} dt.$$

Proposition 5.1 ([14, Proposition 4.4]). *Suppose that ϕ_1 and ϕ_2 satisfy the doubling condition (3.1). For ϕ_1 , define Φ_1^* and Φ_1^{**} by (5.1) and (5.2), respectively. Let $\phi_3 = \phi_2/(\Phi_1^* + \Phi_1^{**})$. If $1 \leq p_2 < p_1 < \infty$ and $p_4 \geq p_1 p_2/(p_1 - p_2)$, then*

$$(5.3) \quad \text{PWM}(\mathcal{L}_{p_1, \phi_1}^{\natural}(\mathbb{R}^n), \mathcal{L}_{p_2, \phi_2}^{\natural}(\mathbb{R}^n)) \supset \mathcal{L}_{p_2, \phi_3}^{\natural}(\mathbb{R}^n) \cap L_{p_4, \phi_2/\phi_1}(\mathbb{R}^n),$$

$$(5.4) \quad \|g\|_{\text{Op}} \leq C(\|g\|_{\mathcal{L}_{p_2, \phi_3}^{\natural}} + \|g\|_{L_{p_4, \phi_2/\phi_1}}),$$

where $\|g\|_{\text{Op}}$ is the operator norm of $g \in \text{PWM}(\mathcal{L}_{p_1, \phi_1}^{\natural}(\mathbb{R}^n), \mathcal{L}_{p_2, \phi_2}^{\natural}(\mathbb{R}^n))$.

Lemma 5.2 ([14, Lemma 3.5]). *Let $1 \leq p < \infty$. Suppose that ϕ satisfies the doubling condition (3.1). Then*

$$(5.5) \quad \mathcal{L}_{p, \phi}^{\natural}(\mathbb{R}^n) \subset L_{p, \Phi^* + \Phi^{**}}(\mathbb{R}^n) \quad \text{and} \quad \|f\|_{L_{p, \Phi^* + \Phi^{**}}} \leq C \|f\|_{\mathcal{L}_{p, \phi}^{\natural}}.$$

Corollary 5.3. *Suppose that ϕ satisfies the doubling condition (3.1). Let $\psi = \phi(\Phi^* + \Phi^{**})$.*

If $1 \leq p_2 < p_1 < \infty$ and $p_4 \geq p_1 p_2/(p_1 - p_2)$, then

$$(5.6) \quad \text{PWM}(\mathcal{L}_{p_1, \phi}^{\natural}(\mathbb{R}^n), \mathcal{L}_{p_2, \psi}^{\natural}(\mathbb{R}^n)) \supset \mathcal{L}_{p_4, \phi}^{\natural}(\mathbb{R}^n),$$

$$(5.7) \quad \|g\|_{\text{Op}} \leq C \|g\|_{\mathcal{L}_{p_4, \phi}^{\natural}},$$

where $\|g\|_{\text{Op}}$ is the operator norm of $g \in \text{PWM}(\mathcal{L}_{p_1, \phi}^{\natural}(\mathbb{R}^n), \mathcal{L}_{p_2, \psi}^{\natural}(\mathbb{R}^n))$. This implies that

$$(5.8) \quad \|fg\|_{\mathcal{L}_{p_2, \psi}^{\natural}} \leq C\|f\|_{\mathcal{L}_{p_1, \phi}^{\natural}} \|g\|_{\mathcal{L}_{p_4, \phi}^{\natural}}.$$

For example, we can take $p_1 = p_4 = 4$ and $p_2 = 2$.

Proof. By Lemma 5.2 we have the inclusion

$$(5.9) \quad \mathcal{L}_{p_2, \phi}^{\natural}(\mathbb{R}^n) \cap L_{p_4, \Phi^* + \Phi^{**}}(\mathbb{R}^n) \supset \mathcal{L}_{p_4, \phi}^{\natural}(\mathbb{R}^n),$$

$$(5.10) \quad \|g\|_{\mathcal{L}_{p_2, \phi}^{\natural}} + \|g\|_{L_{p_4, \Phi^* + \Phi^{**}}} \leq C\|g\|_{\mathcal{L}_{p_4, \phi}^{\natural}}.$$

Then, using Proposition 5.1, we have the conclusion. $\square$

6. SPECIFIC FUNCTION SPACES

We now give the specific function spaces V and W satisfying (1.6), (1.7) and (1.8).

For example, let $p > 2$, $-n/p \leq \alpha_* < 0 < \alpha < 1$, $-n/p \leq \beta < 0$, and

$$(6.1) \quad \phi(x, r) = \begin{cases} r^{\alpha}, & |x| \leq 2, \ 0 < r \leq 2, \\ r^{\beta}, & |x| \leq 2, \ r > 2, \\ r^{\alpha_*}, & |x| > 2, \ 0 < r \leq 2, \\ r^{\beta}, & |x| > 2, \ r > 2, \end{cases} \quad \psi(x, r) = \begin{cases} r^{\alpha}, & |x| \leq 2, \ 0 < r \leq 2, \\ r^{\beta}, & |x| \leq 2, \ r > 2, \\ r^{2\alpha_*}, & |x| > 2, \ 0 < r \leq 2, \\ r^{\beta}, & |x| > 2, \ r > 2, \end{cases}$$

and take

$$W = \mathcal{L}_{p/2, \psi}^{\natural}(\mathbb{R}^n) \quad \text{and} \quad V = \mathcal{L}_{p, \phi}^{\natural}(\mathbb{R}^n),$$

then V and W satisfy (1.6), (1.7) and (1.8) when $n = 3$. We will check these properties in this section.

Firstly, we see that ϕ and ψ satisfy (3.1) and

$$\psi(x, r) = r^{\alpha} \quad \text{for all } B(x, r) \subset B(0, 2).$$

Then, by Proposition 3.4, we have

$$\|f\|_{\text{Lip}_{\alpha}(B(0, 2))} \leq C\|f\|_{\mathcal{L}_{p/2, \psi}},$$

and

$$\|f\|_{\mathcal{L}_{p/2, \psi}^{\natural}} \sim \|f\|_{\mathcal{L}_{p/2, \psi}} + |f(0)|.$$

This shows the property (1.6). Next, the properties (1.7) and (1.8) follows from Propositions 6.1 and 6.2 below, respectively. Therefore, if $f, g \in \mathcal{L}_{p, \phi}^{\natural}(\mathbb{R}^n)$ and $\sigma(fg) = \lim_{r \rightarrow \infty} (fg)_{B(0, r)} = 0$, then

$$|(R_j R_k(fg))(0)| \leq \|R_j R_k(fg)\|_{\mathcal{L}_{p/2, \psi}^{\natural}} \leq C\|fg\|_{\mathcal{L}_{p/2, \psi}^{\natural}} \leq C\|f\|_{\mathcal{L}_{p, \phi}^{\natural}} \|g\|_{\mathcal{L}_{p, \phi}^{\natural}}.$$

Further, let f be α -Lipschitz continuous on $B(0, 2)$ and $|f(x)| \leq C/|x|$ for $|x| \geq 2$. Then $\sigma(f) = 0$ and f is in $\mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$, if p and β satisfy one of the following conditions:

$$\begin{cases} 2 < p < n & \text{and } -1 \leq \beta < 0, \\ p = n & \text{and } -1 < \beta < 0, \\ n < p & \text{and } -n/p \leq \beta < 0. \end{cases}$$

Moreover, if $\alpha_* = \beta/2 = -n/p$ also, then $-n/(p/2) = 2\alpha_* = \beta < 0$ and

$$\|R_j R_k(fg)\|_{\text{Lip}_\alpha(B(0,2))} + \|R_j R_k(fg)\|_{L^{p/2}} \leq C \|R_j R_k(fg)\|_{\mathcal{L}_{p/2,\psi}^{\natural}} \leq C \|f\|_{\mathcal{L}_{p,\phi}^{\natural}} \|g\|_{\mathcal{L}_{p,\phi}^{\natural}},$$

for all $f, g \in \mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$ satisfying $\sigma(fg) = 0$, see Proposition 3.5.

Note that, in the decomposition $u = U + r$ in Definition 1.2, we may assume that U has a compact support in $\mathbb{R}^3$ at fixed t . Then $|r(t, x)| \leq C/|x|$ for large $x \in \mathbb{R}^3$. It is also known that $\nabla u \in L^\infty(\mathbb{R}^3)$ at t , see [10], that is, ∇r is bounded. Hence $\sigma(\partial_3 r_i U_j) = \sigma(r_i \partial_3 U_j) = \sigma(r_i \partial_3 r_j) = 0$ for all i, j .

Proposition 6.1. *Let $p \geq 2$, $-n/p \leq \alpha_* < 0 < \alpha \leq 1$, $-n/p \leq \beta < 0$, and let ϕ and ψ be as (6.1). Then there exists a positive constant C such that, for all $f, g \in \mathcal{L}_{p,\phi}^{\natural}(\mathbb{R}^n)$,*

$$(6.2) \quad \|fg\|_{\mathcal{L}_{p/2,\psi}^{\natural}} \leq C \|f\|_{\mathcal{L}_{p,\phi}^{\natural}} \|g\|_{\mathcal{L}_{p,\phi}^{\natural}}.$$

Proof. For ϕ in (6.1), we have

$$\Phi^*(x, r) = \int_1^{\max(2, |x|, r)} \frac{\phi(0, t)}{t} dt = \int_1^2 t^{\alpha-1} dt + \int_2^{\max(2, |x|, r)} t^{\beta-1} dt \sim 1,$$

and

$$\begin{aligned} 1 + \Phi^{**}(x, r) &= 1 + \int_r^{\max(2, |x|, r)} \frac{\phi(x, t)}{t} dt \\ &= 1 + \begin{cases} \int_r^2 t^{\alpha-1} dt, & |x| \leq 2, 0 < r \leq 2, \\ 0, & |x| \leq 2, r > 2, \\ \int_r^2 t^{\alpha_*-1} dt + \int_2^{|x|} t^{\beta-1} dt, & |x| > 2, 0 < r \leq 2, \\ \int_r^{\max(|x|, r)} t^{\beta-1} dt, & |x| > 2, r > 2, \end{cases} \\ &\sim \begin{cases} 1, & |x| \leq 2, 0 < r \leq 2, \\ r^{\alpha_*}, & |x| > 2, 0 < r \leq 2, \\ 1, & r > 2. \end{cases} \end{aligned}$$

Hence

$$\phi(x, r)(\Phi^*(x, r) + \Phi^{**}(x, r)) \sim \psi(x, r) = \begin{cases} r^\alpha, & |x| \leq 2, 0 < r \leq 2, \\ r^{2\alpha_*}, & |x| > 2, 0 < r \leq 2, \\ r^\beta, & r > 2. \end{cases}$$

Then, using Corollary 5.3, we have the conclusion. $\square$

Proposition 6.2. *Let $q > 1$, $-n/q \leq \delta < 0 < \alpha < 1$, $-n/q \leq \beta < 0$, and*

$$\psi(x, r) = \begin{cases} r^\alpha, & |x| \leq 2, \ 0 < r \leq 2, \\ r^\beta, & |x| \leq 2, \ r > 2, \\ r^\delta, & |x| > 2, \ 0 < r \leq 2, \\ r^\beta, & |x| > 2, \ r > 2. \end{cases}$$

Then the Riesz transforms $\tilde{R}_j$, $j = 1, 2, \dots, n$, are bounded on $\mathcal{L}_{q,\psi}(\mathbb{R}^n)$ and on $\mathcal{L}_{q,\psi}^{\natural}(\mathbb{R}^n)$. That is, there exists a positive constant C such that, for all $f \in \mathcal{L}_{q,\psi}(\mathbb{R}^n)$,

$$\|\tilde{R}_j f\|_{\mathcal{L}_{q,\psi}} \leq C \|f\|_{\mathcal{L}_{q,\psi}}, \quad \|\tilde{R}_j f\|_{\mathcal{L}_{q,\psi}^{\natural}} \leq C \|f\|_{\mathcal{L}_{q,\psi}^{\natural}}, \quad j = 1, 2, \dots, n.$$

Moreover, if $f \in \mathcal{L}_{q,\psi}^{\natural}(\mathbb{R}^n)$ and $\sigma(f) = \lim_{r \rightarrow \infty} f_{B(0,r)} = 0$, then the Riesz transforms $R_j f$, $j = 1, 2, \dots, n$, are well defined, $\sigma(R_j f) = \lim_{r \rightarrow \infty} (R_j f)_{B(0,r)} = 0$, and

$$\|R_j f\|_{\mathcal{L}_{q,\psi}^{\natural}} \leq C \|f\|_{\mathcal{L}_{q,\psi}^{\natural}}, \quad j = 1, 2, \dots, n.$$

Proof. We see that ψ satisfies (3.1) and

$$r \int_r^\infty \frac{\psi(x, t)}{t^2} dt \leq A \psi(x, r),$$

for all $x \in \mathbb{R}^n$ and $r > 0$. Then we have the boundedness of $\tilde{R}_j$ on $\mathcal{L}_{q,\psi}(\mathbb{R}^n)$ and on $\mathcal{L}_{q,\psi}^{\natural}(\mathbb{R}^n)$. Let

$$\tilde{\psi}(x, r) = \tilde{\psi}(r) = \begin{cases} r^\delta, & 0 < r \leq 2, \\ r^\beta, & r > 2. \end{cases}$$

Then $\tilde{\psi}$ satisfies (3.1), (3.2), (3.5) and $\psi \leq \tilde{\psi}$. Therefore, by Theorem 4.4, we have the conclusion. $\square$

ACKNOWLEDGMENTS

The first author was partially supported by Grant-in-Aid for Scientific Research (C), No. 24540159, Japan Society for the Promotion of Science. The second author was partially supported by Grant-in-Aid for Young Scientists (B), No. 25870004, Japan Society for the Promotion of Science.

REFERENCES

- [1] J. T. Beale, T. Kato, and A. Majda, *Remarks on the breakdown of smooth solutions for the 3-D Euler equations*, Comm. Math. Phys., 94 (1984), 61–66.
- [2] L. Caffarelli, R. Kohn and L. Nirenberg, *Partial regularity of suitable weak solutions of the Navier-Stokes equations*, Comm. Pure Appl. Math., 35 (1982), 771–831.
- [3] D. Chae, *Local existence and blow-up criterion for the Euler equations in the Besov spaces*, Asymp. Anal. 38 (2004), 339–358.
- [4] C-H. Chan and T. Yoneda, *On possible isolated blow-up phenomena and regularity criterion of the 3D Navier-Stokes equation along the streamlines*, Methods Appl. Anal., 19 (2012), 211–242.
- [5] P. Constantin and C. Fefferman, *Direction of vorticity and the problem of global regularity for the Navier-Stokes equations*, Indiana Univ. Math. J., 42 (1993), 775–789.

- [6] P. Constantin, C. Fefferman and A. Majda, *Geometric constraints on potential singularity formulation in the 3D Euler equations*, Comm. Partial Differential Equations, 21 (1996), 559–571.
- [7] J. Deng, T.-Y. Hou and X. Yu, *Improved geometric conditions for non-blowup of the 3D incompressible Euler equation*. Comm. Partial Differential Equations, 31 (2006), 293–306.
- [8] C. Fefferman, *Existence and smoothness of the Navier-Stokes equation*, The millennium prize problems, 57–67, Clay Math. Inst., Cambridge, MA, 2006.
- [9] Y. Giga, *Solutions for semilinear parabolic equations in L^p and regularity of weak solutions of the Navier-Stokes system*, J. Differential Equations, 62 (1986), 186–212.
- [10] Y. Giga, K. Inui and S. Matsui, *On the Cauchy problem for the Navier-Stokes equations with nondecaying initial data*. Advances in fluid dynamics, 27–68, Quad. Mat., 4, Dept. Math., Seconda Univ. Napoli, Caserta, (1999).
- [11] H. Kozono and Y. Taniuchi, *Bilinear estimates in BMO and the Navier-Stokes equations*, Math. Z., 235 (2000), 173–194.
- [12] E. Nakai, *Pointwise multipliers for functions of weighted bounded mean oscillation*, Studia Math., 105 (1993), 105–119.
- [13] E. Nakai, *Hardy-Littlewood maximal operator, singular integral operators and the Riesz potentials on generalized Morrey spaces*, Math. Nachr., 166 (1994), 95–103.
- [14] E. Nakai, *Pointwise multipliers on weighted BMO spaces*, Studia Math., 125 (1997), 35–56.
- [15] E. Nakai, *The Campanato, Morrey and Hölder spaces on spaces of homogeneous type*, Studia Math., 176 (2006), 1–19.
- [16] E. Nakai, *A generalization of Hardy spaces H^p by using atoms*, Acta Math. Sinica, 24 (2008), 1243–1268.
- [17] E. Nakai, *Singular and fractional integral operators on Campanato spaces with variable growth conditions*, Rev. Mat. Complut., 23 (2010), 355–381.
- [18] E. Nakai and Y. Sawano, *Hardy spaces with variable exponents and generalized Campanato spaces*, J. Funct. Anal., 262 (2012), 3665–3748.
- [19] Eiichi Nakai and Kôzô Yabuta, *Pointwise multipliers for functions of bounded mean oscillation*, J. Math. Soc. Japan, 37 (1985), 207–218.
- [20] E. Nakai and T. Yoneda, *Bilinear estimates in dyadic BMO and the Navier-Stokes equations*. J. Math. Soc. Japan, 64 (2012), 399–422.

DEPARTMENT OF MATHEMATICS, IBARAKI UNIVERSITY, MITO, IBARAKI 310-8512, JAPAN
E-mail address: enakai@mx.ibaraki.ac.jp

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, MEGURO-KU, TOKYO 152-8551, JAPAN
E-mail address: yoneda@math.titech.ac.jp